



History:

Received: 25 July 2020

Accepted: Sep. 01, 2020

Published: Sep. 17, 2020

Collection year: 2020

Confirmation of publication:

Published

Identifiers and Pagination:

Year: 2020

Volume: 12

First Page: 39

Last Page: 56

Publisher ID:

19204159.12.19

DOI:

<http://dx.doi.org/10.21065/19204159.12.19>

19204159.12.19

Corresponding author:

Muhammad Modassar Ali

Nawaz Ranjha Institute of

Food Science and Nutrition,

University of Sargodha,

Sargodha – Pakistan

E:modassarranjha@gmail.com

m

Citation:

Maria B. et al., A Critical
Review on Alpha Tocopherol:
Sources, RDA And Health
Benefits Vol. 12: 2020. p. 39-
56

Review article

A CRITICAL REVIEW ON ALPHA TOCOPHEROL: SOURCES, RDA AND HEALTH BENEFITS.

Maria Batool¹, Samina Kauser², Hafiz Rehan Nadeem³, Rashida Perveen⁴, Shafeeqa Irfan², Ayesha Siddiqua³, Bakhtawar Shafique², Syeda Mahvish Zahra², Muhammad Waseem³, Waseem Khalid⁵, Muhammad Modassar Ali Nawaz Ranjha².

1. University Institute of Diet and Nutritional Sciences, University of Lahore, Gujrat – Pakistan
2. Institute of Food Science and Nutrition, University of Sargodha, Sargodha – Pakistan
3. Institute of Food Science and Nutrition, Bahaddin Zakariya University, Multan – Pakistan
4. Department of Allied Health Sciences, Superior College, University Campus Lahore, Pakistan
5. Institute of Home and Food Sciences, Government College University, Faisalabad – Pakistan

Abstract:

Virtually 100 years before α -tocopherol (universally known as a “Vitamin-E”) remained revealed. It was essential component to preclude fetal recommencement in expectant. In 1940s, Vitamin E existed recognized as the crucial part that plays in deactivating and extinguishing the free radicals that is present inside the diverse compartments of muscles. There are the eight diverse types of bits interconnected to Alpha-tocopherol (Vitamin E) are founded in human food or diet that is basically manufactured from plants. In overall, the quantity of α -tocopherol (as an Vitamin E) provided by a standard nourishment assortments from 6 to 13 mg per day (1 IU = 0.67 mg). In record states, the predictable mediocre consumption is lower than 10 mg per day. Lipoproteins that are act as an antioxidant are the form of vitamin E is produced in different parts of the body especially in Liver organ where it plays as defender of definite fatty acids and proteins that is existent in the tissue from free activists. It alleviates the cellular casings and it also enhanced the enablement of the immune reaction throughout the infections. It has been reported that the Vitamin E is used to be protecting the various types of diseases like it save the colon infection, prostate malignance, certain cardiac syndromes, ischemia, flume, inflammation and nervous infections.

Keywords: α -tocopherol, Vitamin-E, Health benefits,

ALPHA TOCOPHEROLS

All plants synthesize α -tocopherols (Vitamin E), which are deliberated as lipophilic antioxidants. They stow lipid peroxyl radicals during intense antioxidant activities (Massacci *et al.*, 2008). The function of α -Tocopherols is to deliver a defense environment to assorted structures, lipids and additional membranes (Nordberg and Arnér, 2001). On interrelating with acyl groups of lipids, α -tocopherols scavenge many lipid soluble and reactive oxygen species (ROS) compounds produced during stress of oxidation process (Munné-Bosch, 2005). Drought-stressed plants have been recounted to synthesize α -tocopherols (Martinez *et al.*, 2007). Purpose of oxidative stress is actually to activate the genes countenances that are accountable for tocopherols synthesis in shrubberies (Falk *et al.*, 2002). It was testified and reported that, vigna plants have an enhanced number of tocopherols (Manivannan *et al.*, 2007). Zhang and Schmidt (2000) described a twice intensification in α -tocopherol in lawn mowers under water stress. Synthesis of α -tocopherol is under genetic control activated by stress produced by oxidation in plants (Gong *et al.*, 2012).

There are 08 naturally happening vitamin E having the following biochemical arrangements (gamma, alpha, beta, delta tocopherols and gamma, beta, delta and alpha toco-trienols) having a fluctuating stage of genetic commotion (Shils and Shike, 2006). α -tocopherol is the one that is considered to encounter body requirements of human beings. Meditation of serum of vitamin E

Funding:

The authors received no direct funding for this research.

Competing Interests:

The authors declare no competing interests

Additional information is available at the end of the article.

(alpha-tocopherol) mainly hangs on the liver, which engrosses numerous nutrients from small intestine and engrosses the nutrient. The liver preferentially releases alpha-tocopherol only through the hepatic alpha-tocopherol transfer protein (Shils and Shike, 2006). Liver removes and metabolizes an additional form of vitamin E (Traber, 2007). Consequently, cell concentrations and blood of additional forms of vitamin E are inferior to alpha-tocopherol and have been the topic of fewer investigations (Dietrich *et al.*, 2006).

Sources of Vitamin E

Products from natural sources are being used from centuries (Ranjha et al., 2020a; Ranjha et al., 2020b). Numerous foods are enriched with vitamin E. Vegetable oils, nuts and seeds are the superlative foundations of α -tocopherol and are found in momentous volume in fortified cereals and green leafy vegetables (Grasp Table 1 & 2 for a detail of quantity of α -tocopherol in diverse foods and RDAs) (Canales *et al.*, 2019).

Table 1: Best Sources of Alpha-Tocopherol

Food	Per serving (mg)	Daily value (%)
Spinach, Raw (01 Cup)	0.6	3
Tomato, Fresh (01)	0.7	4
Mango, Sliced (½ Cup)	0.7	4
Kiwifruit (01 Medium)	1.1	6
Oil of Soybean (01 Tbs)	1.1	6
Broccoli, Boiled, Chopped (½ Cup)	1.2	6
Spinach, Cooked (½ Cup)	1.9	10
Oil of Corn (01 Tbs)	1.9	10
Peanuts, Dry Roasted (01 Oz)	2.2	11
Butter of Peanut (02 Tbs)	2.9	15
Parched Roasted Hazelnuts (01 Oz)	4.3	22
Oil of Safflower (01 Tbs)	4.6	25
Oil of Sunflower (01 Tbs)	5.6	28
Dry Roasted Almonds (01 Oz)	6.8	34
Dry Roasted Seeds of Sunflower, (01 Ounce)	7.4	37
Oil of Wheat Germ (01 Tbs)	20.3	100

Maximum vitamins E in diets of Americans are in form of gamma-tocopherols. Vegetable oils, corn, canola and soy have an extended capacity of vitamin E (Dietrich *et al.*, 2006).

Table 2: RDAs for Vitamin E (α -Tocopherol) with respect to age and gender (DRI, 2005)

Age	Women	Men	Gestation	Lactation
0 – 6 Months*	4 mg 6 (IU)	4 mg 6 (IU)	-	-
7 – 12 Months*	5 mg 7.5 (IU)	5 mg 7.5 (IU)	-	-
1 – 3 Years*	6 mg 9 (IU)	6 mg 9 (IU)	-	-
4 – 8 Years*	7 mg 10.4 (IU)	7 mg 10.4 (IU)	-	-

9 – 13 Years*	11 mg 16.4 (IU)	11 mg 16.4 (IU)	-	-
14 + Years*	15 mg 22.4 (IU)	15 mg 22.4 (IU)	15 mg 22.4 (IU)	19 mg 28.4 (IU)

Peroxidation of lipids is a destructive and free radical scavenging process which is accountable for development of aromas in fats, foods and oils (Babu, 2011). In addition, oxidation of unsaturated fatty acids from bio-mass actually constructs pathological and functional aberrations and variations (Schaich, 2006). Tocotrienols and tocopherols are better documented for their operative cessation of peroxidation of lipids in living cells and foods (Burton & Traber, 1990). Synthesis of vitamin E takes place in plants only. Consequently, it is regarded as an imperative nutrient for both animals and humans (Tegischer *et al.*, 2002).

Isomers of tocopherols are chain-contravention anti-oxidants. The anti-oxidative commotion of tocopherols is accompanying to scavenge free radicals of un-saturated lipids (Chepda *et al.*, 2001; Fujisawa *et al.*, 2004). α -Tocopherols are an utmost organically abundant and active state of vitamin E (in vivo) which competently transferences a H^+ atom to a lipid free radical, such as carbon centered, alkoxyl and peroxy radicals, bountiful consistent non-radical artifact of an α -tocopheroxyl radical and lipid. After development, α -tocopherol radicals react with a second free radical or with each other to form a non-radical product. Therefore, each α -tocopherol molecule consumes 2 free lipid radicals and terminates the free radical chain reaction. To elucidate the mechanism of tocopherol free radical modification, the reaction products of tocopherols with lipid free radicals have been investigated.

Health Assistances

In cellular anti-oxidant protection mechanism, vitamin E is regarded as a most important component that is lipid-soluble and this is completely attained from food. Due to an anti-oxidant action, vitamin E plays frequent imperative characters inside the body. Process of oxidation is associated with numerous conceivable diseases and circumstances, counting cataract, arthritis, aging and cancer. Vitamin E has a capability to preclude from atherosclerosis and unnecessary upsurge in platelets. Furthermore, it may help to decrease the prostaglandin production like thromboxane that persuades aggregation of platelets (Rizvi *et al.*, 2014).

Antioxidant Activity of α -tocopherol

Ohm *et al.*, 2005 accompanied a study to evaluate the antioxidant action of α -tocopherol upon oxidation of lipids. According to them, α -tocopherols are recognized as influential antioxidants. Objective of that study (in vitro) was to inspect the dosage-based reserve of formation of oxidation product instigated by α -tocopherol and also to guesstimate potential of peak anti-oxidant action of α -tocopherol at diverse phases of oxidation of lipids. In rapeseed oil tri-glycerides (purified rapeseed oil\ ROTGs), α -tocopherol was added in some meditations fluctuating from 25 – 1500 mmol/ kg. The antioxidant action of α -tocopherol augmented fit for a meditation of 100 mmol/ kg (purified rapeseed oil). A meditation of 125 mmol of α -tocopherol/ kg (purified rapeseed oil) didn't show an enhanced anti-oxidant outcome. The development of subordinate oxidation products (volatile) tracked analogous trend and as a result, an extreme anti-oxidant consequence was seen for 100 mmol α -tocopherol/ kg. Additionally, meditations among 250 and 1500 mmol α -tocopherol / kg (purified rapeseed oil) evidently triggered the augmented production of hydro-peroxides through period of induction. As interrelated with control group, entire tried mediations of α -tocopherols occasioned in reduced hydro-peroxide creation and no pro-oxidative properties were pragmatic.

Traber and Atkinson, (2007) presented a theory that all annotations concerning the mechanism of in vivo initiation of α -tocopherol stem from its role as a lipid soluble anti-oxidant. Purpose of that study was to designate an evidence for in-vivo effectiveness of alpha-tocopherol. The role of alpha-tocopherol as an anti-oxidant is to uphold the legitimacy of long-chain fatty acids (unsaturated) in superficial surface (cell membrane) and thus sustaining a normal genetic commotion. It means, such bioactive lipids are necessary beckoning particles which alters in their quantities or forfeiture because of oxidation and recognized as the chief events in cells which are retorted by other cells. Innumerable pathways which have been defined by another in order to be lowered than alpha-tocopherol parameter seem somewhat to be reliant on the oxidative pressure of tissue under question or cells. Furthermore, it seemed improbable that those pathways were exactly beneath the regulation of alpha-tocopherol assumed that numerous anti-oxidants excluding various oxidative stressors and alpha-tocopherol might deploy their retorts. Consequently, entire scope and variations of alpha-tocopherol's organic commotion could be understandable in light of the membrane qualities (lipid domains, phase separation and fluidity) and defense of poly-unsaturated fatty acids which bring about by poly-unsaturated fatty acids.

Dandapat *et al.* (2000) conducted a study in order to regulate an outcome of added α -referncetocopherol in 200, 400 and 600 mg/ kg feed upon anti-oxidant defense system and lipid peroxidation (LPX) in gills and pancreatic cells of the shrimp (*Macrobrachium rosenbergii*) According to outcomes of this study, it was showed that α -tocopherol repressed an augmented lipid peroxidation (LPX) in gills as compared to hepatopancreas. To the entire 03 supplemented regime, superoxide dismutase (SOD, EC 1.15.1.1) activity was reduced expressively in gills. Reduction in pancreatic cells were detected just in retort to advanced dosages of α -tocopherol i. e. 400 and 600 mg/ kg feed. Enzymatic action was not abridged pointedly in pancreatic cells but gills. The activity of glutathione peroxidase (GPx) (EC 1.11.1.9) remained uninfluenced in gills. There's highest quantity in the pancreatic cells by the consumption of α -tocopherol. Glutathione-disulfide reductase (GSR) action remained same in pancreatic cells but declined in gills. A considerable augmentation in the quantity of glutathione (GSH) content was noticed in both the pancreatic cells and gills. While, content of vitamin C (ascorbic acid) raised suggestively in pancreatic cells and remained unchanged in gills. So, conclusions of this enquiry proposed that vitamin-E has a capability to reduce level of LPX and also might control anti-oxidant defense system in pancreatic cells and gills, rejoinder was tissue specific exceedingly. Further, it was detected that the consumption of high dosage of α -tocopherol (i. e. 600 mg/ kg feed) is not capable to reduce considerable extra-defense in gills and pancreatic cells.

Tucker and Townsend, (2005) reviewed the health outcomes of α -tocopherol in therapy, and prevention from innumerable human diseases. α -tocopherol is recognized as an imperative out of the 08 types of tocopherol (Vitamin E). α -tocopherol is an utmost effective anti-oxidant (fat-soluble) acknowledged on landscape. It was a myth in past those α -tocopherol acts only as a lipid peroxy radicals' scavenger. Unambiguously, with oxidized low-density lipoprotein (oxLDL), thus helping as an imperative anti-oxidant for inhibition of arterial hardens (atherosclerosis); now days, evidence-based studies claimed many of the adventitious effects of α -tocopherol (other than anti-oxidant action) which comprise gene regulatory, pro-oxidant, cell signaling belongings. Periods of preclinical and clinical investigations have extended our consideration regarding anti-oxidant action of α -tocopherol and its efficacy in multiple and enduring oxidative stress-tempted pathogenesis. Outcomes of this investigation has publicized even though mixed and auspicious assessments on effectiveness of anti-oxidant action of α -tocopherol in handling and deterrence of Alzheimer's disease, cancer and cardiovascular disorders. It was advocated that more investigations to expose

systemic and cellular mechanisms might assist in supervision of apposite medical dealing approaches by utilizing α -tocopherol transversely in an assorted populace of aged entities.

Anti-Inflammatory Commotion of α -Tocopherol

Grammas *et al.* (2004) scrutinized the anti-inflammatory possessions of metabolites of tocopherol (c-tocopherol, α -tocopherol) and its association with 2,7,8-trimethyl-2-(b-carboxyethyl)-6-hydroxychroman (c-CEHC) and 2,5,7,8-tetramethyl-2-(b-carboxyethyl)-6-hydroxychroman (α -CEHC) and in a distinct culture system of cells. An aortic cell of endothelium of mice and cultures of microglia was treated with tumor necrosis factor (cachectin, cachexin, or TNF), prostaglandin E₂ (PGE₂) and bacterial lipopolysaccharide (LPS) and nitrite was analyzed. TNF α -stimulated nitrite production was repressed by α -CEHC in both types of cells, while inhibition of both CEHC derivatives and LPS-stimulated microglial nitrite efflux resulted. Both c-CEHC and α -CEHC repressed the production microglia PGE₂, but neither c- nor α -tocopherol proved operative in inhibition of inflammatory processes of stimulated cytokine. According to the consequences, it was proved that anti-inflammatory properties of tocopherols were extremely endpoint-dependent, stimulated and specific cell type.

Atherogenicity action and Other Cardiac Outcomes

Oxidative variation of low-density lipoprotein (LDL) is supposed to prove a significant character in development of atherosclerosis (Parthasarathy *et al.*, 2010). Low-density lipoprotein (LDL-Oxidized) is quickly reserved by microorganisms and grounds accumulation of lipids and fluffs the formation of cell. It is recognized as a biochemical for T-lymphocytes and monocytes, that enhance the production of growth factor and inflammatory cytokines by which arterial response and cellular activity enhances (Serin *et al.*, 2007). The reputation of lipoprotein oxidation in thermogenesis is actually a consequence of measurement of rapid thickening of the carotid artery wall and lipoprotein oxidation in adult a human being (Nadeem *et al.*, 2012). In previous few years, it has become a topic of interest that diets enriched with vitamins (anti-oxidants) have a capability to condense the menace of atherosclerosis by averting low-density lipoprotein (LDL) by modification in oxidative processes. Inhibition of lipid peroxidation (ex-vivo) was shown by supplements encompassing antioxidant compounds, carotene, vitamin C and vitamin E (H, 2010). In numerous observational researches, it has been stated that advanced consumptions and blood meditations of vitamins were linked with an abridged risk of coronary artery disease and stroke (Ryan *et al.*, 2007). Limited investigations have shown a connotation among early stages of atherosclerosis and utilization of α -tocopherols (vitamin E anti-oxidants). Finding of atherosclerosis in extra-corporeal carotid arteries is possible with an aid of dual ultrasound and it is a sensitive, specific and noninvasive technique. An atherosclerotic plaque is considered responsible for augmented intima-carotid thickness in the arterial system and envisages stroke and myocardial infarction in normal healthy individuals (Raitakari *et al.*, 2003). Cardiovascular diseases (CVDs) result due to dietary factors (Alissa and Ferns, 2017) so; optimization of nutritional parameters must be well thought-out as a significant routine interference in managing of existing cardiovascular diseases (CVDs) and also its anticipation (Mead *et al.*, 2006).

It has been evidently proved and verified that diet plays an imperative role in maintaining good health, counting explanations that the frequency of explicit illnesses diverges extensively across nations. In such type of problematic situations, it is highly mandatory to alter the diet, which is recommended as safe (Alissa and Ferns, 2017). α -tocopherols has a potential to treat and lessening the risk of cancer (Alissa and Ferns, 2017). Many evidences of cardiovascular diseases (CVDs) and diet have testified relations of danger with definite nutrients instead of whole nutriments (Satija *et al.*,

2015). However, innumerable investigations stated the character of complete nutritional outlines in expecting cardiovascular diseases (CVDs) (Denova-Gutiérrez *et al.*, 2016).

Stroke and coronary heart disease are the 02 major ailments caused due to atherosclerosis that is an enduring provocative illness (Yoshida, 2018). The characterization of atherosclerosis is finalized\ diagnosed by the presence of cholesterol accretion and intimal plaques in wall of arteries (Zhu *et al.*, 2018). An inequity among radical scavenging systems (anti-oxidant protection) and radical creation (nitrogen species development or reactive O₂) is thought to be a root etiology of oxidative tension (Refa *et al.*, 2019). Furthermore, when atherosclerosis is in developmental stages, it happens due to many prominent and fatal risk factors comprising bacterial infection, a genetic predisposition, smoking, stress, a high level of C-reactive protein (CRP), dyslipidemia, hypercholesterolemia, obesity, insulin resistance, diabetes mellitus, hypertension, vascular wall inflammation, an immunological disorder and alcohol consumption (Vaccarezza *et al.*, 2018).

In the cure and treatment of atherosclerosis, conventional, nutritional and herbal medication are preferred due to their cost effectiveness and fewer adverse effects as compared to allopathic drugs which are hazardous due to its maximum adverse effects (Chen *et al.*, 2019). Atherosclerotic development mainly caused due to oxidized low-density lipoproteins (LDL) and reactive oxygen species (ROS) and it is well thought that the intake\ consumption of anti-oxidants might possess a high and effective healing approach to avert the growth of atherosclerosis (Vasiliki *et al.*, 2019). Mediation of per-oxidation take place just due to the presence of pro-oxidant substances which either are non-radical and free radical species. Reactive nitrogen species (RNS) and reactive oxygen species (ROS) are the 02 main classifications. Nitrogen dioxide, Nitric oxide, Peroxyl, Hydroperoxyl, Hydroxyl and Superoxide are recognized as the primary responsive and permitted radical species (Jargin, 2018). Nitrite, peroxyntirite and hydrogen peroxide are the chief responsive non-radical class (Jain, 2019). Mitochondrion is the main source of reactive oxygen species (ROS) inside the cell (Liu *et al.*, 2018). In addition, uncoupled endothelial nitric oxide synthetase (eNOS), nitric oxide synthase (NOS), myeloperoxidase (MPO), cyclooxygenase (COX), lipoxygenase (LOX) and xanthine oxidase (XO) are some of the oxidizing enzymes that ensued in endothelial cells, vascular smooth muscle cells and macrophages. Low density lipoproteins (LDL), production of RNS\ ROS and oxidation of smooth muscle cell proliferation caused due to these enzymes (Sun *et al.*, 2019).

Irritable oxygen species (ROS) are proficient to mutilation functions of cells and bio-molecules like carbohydrates, proteins and lipids and might be started e. g., hyperglyceridemia and high blood sugar resulting in LDL oxidation and lipid peroxidation (Graves and Bauer, 2018). Nevertheless, a slightly propensity to pursue harm is be contingent not solitary on the spot and the amount of reactive oxygen species (ROS) formed but also on the compensatory manufacturer's rejoinder (Graves and Bauer, 2018). The process of lipoprotein's oxidation is counted as the 1st stage in progression of atherosclerosis having very toxic and deleterious possessions on cells of endothelium (Kim, 2018). Afterward, endothelial dysfunction is caused due to Ox-LDL which might tempt the formation of foam cell, propagation of smooth muscle cells and countenance of adhesion molecules (Hu *et al.*, 2018). Other multifaceted procedures and inflammation are involved in atherosclerosis and this disease might start in early age (Hu *et al.*, 2018). There are 03 phases of development of atherosclerosis i. e. atherosclerotic plaque and before this; the induction of atheroma and a fatty streak formation are resulted causing fatal stage called athero-thrombosis (Ganapathi *et al.*, 2018). Accumulation of lipoproteins, particularly the low-density lipoprotein-cholesterol (LDL-C) inside arteries and some peril influences incline to enable the dispersion of low-density lipoprotein-cholesterol (LDL-C) inside the intima. LDL-C is converted into Ox-LDL just

because of RNS and ROS which leftovers in intima (Malekmohammad *et al.*, 2019).

Oxidized lipids and cytokines are accountable for the initiation of endothelial cells. On the surface of endothelium, cytokines like interferon gamma (IFN- γ), -6 (IL-1, IL-4, IL-6), interleukin-1,-4 and tumor necrosis factor (TNF- α) make countenance of adhesion molecules, monocyte and leukocyte, particularly E-selectin, intercellular adhesion molecule-1 (ICAM) and vascular cell adhesion molecule-1 (VCAM) (Hijmans *et al.*, 2019). By the accumulation of these adhesion molecules, T-lymphocytes and monocytes are generated in the intima of vascular wall (Hijmans *et al.*, 2019). From media to intima, some explicit chemokines are capable to cause migration of smooth muscle cells resulting proliferation of cells. Chemotactic proteins like IL-8, macrophage colony-stimulating factor (M-CSF) and monocyte chemotactic protein-1 (MCP-1) are differentiated through monocytes in sub-endothelial space by macrophages (Brandsma *et al.*, 2019). In foam cell establishment phase, the mono-nuclear phagocytosis takes place and their macrophages identify and approve molecules of Ox-LDL by an aid of scavenger receptors (SRs).

Lastly, the progression of greasy stripes is generated inside the walls of arteries due to the formation of foam cells. Apoptosis (cell death) happened through evolving lesion (atherosclerotic). Due to apoptotic foam cell death, a necrotic core is generated which works as a yard for lipids and cellular debris (Silva *et al.*, 2019). From media – intima, the induction of extracellular matrix production in atheroma formation and few layers are generated due to which a stringy plaque cap (atherosclerotic) is formed through relocation of cells of smooth muscle. T-lymphocytes, macrophages, smooth muscle cells (SMC) and collagen-rich fiber tissues are the parts of cap of atherosclerotic plaque. Lessened flow of blood in vessels and tissue macrophages cause's athermanous lesions (Singh *et al.*, 2018) and formation of atherosclerotic plaque is arbitrated by mechanisms of fibrous cap. TNF and metalloproteinase are secreted by T-lymphocytes and macrophages in the boundaries of an emerging plaque in order to constrain synthesis of collagen in SMC and also lyse extracellular matrix of fibrous cap, correspondingly. Obliteration of fibrous cap takes place due to the breakdown of extra-cellular matrix and thrombogenic contents and coagulation process starts here due to thrombus formation, the adhesion of platelets and blood clot formation (Malekmohammad *et al.*, 2019). Figure 1 portrays the chief events of atherosclerosis.




Figure 1: Events of Atherosclerosis (Adapted from; Malekmohammad *et al.*, 2019)

Alzheimer's syndrome

Evidence recommends that the vitamins or medications that originate an upsurge in quantities of brain catecholamines and defend in contradiction of oxidative damage might deliberate the development of Alzheimer's disease by condensing nerve damage. A double-blind multicenter, placebo-controlled and randomized trial was conducted in patients of Alzheimer's disease. 341 patients received alpha-tocopherol (vitamin E, 2000 IU daily) and discerning mono-amine oxidase blockers (10 mg/ day). Regardless of an arbitrary project, the score at base-line regarding to mini-mental state inspection was much advanced in placebo cluster than in additional 03 groups and that variable was extremely prognostic of principal consequence ($P < 0.001$). In an unadjusted scrutinization, there was no statistically momentous variance in results between 04 clusters. But, the base-line score on the mini-mental state inspection, a noteworthy interruptions in time to the main consequence for patients treated with combination therapy (585 days, $P = 0.049$), alpha-tocopherol (670 days, $P = 0.001$), placebo group (440 days) and selegiline (median time, 655 days; $P = 0.012$), as equated with the patients with abstemiously unadorned damage from Alzheimer's disease (Windisch, 2000).

The progression of Alzheimer's disease (AD), is particularly progressing rapidly in emerging republics and also became an alarming condition for older as well as younger age individuals. Oxidative stress is actually caused due to fluctuated routine alterations that has a capability to cause Alzheimer's disease (AD) and its probability of incidence. Delay would happen in the cure due to improper investigation aids and assistances. For now, only management can be done to relive signs and symptoms. Medications relating to this group of anti-lipid, anti-inflammatory, antioxidants, beta-seccasase, MAO and AChE may exert as a moral treatment in order to get enhanced effectiveness. Many allopathic drugs have been functioned by solitary mechanism. Plants have been operated in multiple ways which inclined to release regressive betterment because of an assorted mode of plants therapy. It was reviewed that current beneficial procedures accessible in the management of Alzheimer's disease (AD). Consequently, no fair treatment is available for the treatment of AD because of the foremost permanent deteriorating brain cells. Complimentary treatments like herbal medications have a tendency to postpone and stop the development of Alzheimer's disease (AD). Current medical information is also recommended. Though, as more and more lead molecules are under clinical investigation, we hope to have unique molecules that can prevent the progression of AD by using the rapid mechanism that is likely to be marketed in the near future (Sivaraman et al., 2019).

Intellectual diminishing

A health study related to women was conducted that was placebo-controlled, double-blind and randomized probationary of vitamin E supplementation *i. e.* 600 IU [α -tocopherol acetate] on different days which started among 39,876 vigorous females of United States (US) in between 1992 and 1995. 6377 females of ≥ 65 years older partaken in that intellectual study. 03 valuations of category fluency, verbal memory and general cognition were directed by an aid of phone at 02 years interlude. A worldwide amalgamated groove is close to recital was the principal conclusion that was applied on all assessments. No variances were noted in worldwide groove among placebo and vitamin E cluster at initial valuation. In the initial phase valuation was done after 5.6 years having (95 % confidence interval [CI], -0.04 – 0.03, mean difference – 0.01) and ended at former valuation at 9.6 years (-0.04 – 0.04, mean difference, 0.00; 95 % CI). Alteration time for mean intellectual capability was not similar in placebo group but similar in vitamin E group ($P = .16$, -0.01 to 0.05, 95 % CI and mean transformation in change = 0.02). The comparative hazard of considerable failure in

worldwide groove in placebo group compared with the vitamin E group was 0.92 (0.77 – 1.10, 95 % CI). It was then proved by this observation analyses that supplementation of vitamin E has no significant role on cognitive behavior (Kang *et al.*, 2006).

Vitamin E is recognized as the valuable and much imperative anti-oxidant in brain in the form of α -tocopherol. Just because of bulky quantities in brain of its carrier α -TTP (α -tocopherol transfer protein) whose purpose includes allocating and regulating levels of vitamin E in numerous cells (Lim and Traber, 2007; Ulatowski and Manor, 2015). Important insights into the function of brain accentuate those carriers of the human mutation in α -TTP gene development advanced and extremely low levels of vitamin E, loss of proliferation, clavicular dysfunction and spino-cerebral ataxia (Doria-Lamba *et al.*, 2006). In encephalon, α -TTP expression is highest in the cerebellum, especially in astrocytes, which supply vitamin E to neighboring neurons. Importantly, α -TTP expression is increased in the brain of patients with neurodegenerative diseases (Ulatowski *et al.*, 2012). So, it has been clinched that vitamin E has an operational character in neuro-protection due to its anti-oxidant action.

Hemodialysis Patients

Augmented mortality and cardiovascular complications are caused due to oxidative stress, particularly in patients facing end-stage renal disease. Though, observational information from the general population have shown that dietary antioxidant intake is associated with reduced cardiovascular morbidity and mortality, most clinical intervention trials have failed to support this relationship. This may be the result of not using an effective antioxidant dose or failing to evaluate patients with high oxidative stress. A study in hemodialysis patients reported that 800 IU / day alpha-tocopherol reduces cardiovascular endpoints. Timely study of recent dose response studies in patients with hypertension and 1,600 IU per day alpha-tocopherol is an optimal dose. Such dose response data are not available for hemodialysis patients. Therefore, the purpose of this study was to evaluate the effect of different doses of oral alpha-tocopherol on oxidative stress in hemodialysis patients with high oxidative stress and the time needed to achieve this effect. The study consisted of a time period and then a dose response study. In a periodic study of 20 hemodialysis patients with high oxidative stress, 1600 IU per day (RRR) alpha-tocopherol for 20 weeks or placebo. Blood was collected every 2 weeks and analyzed for oxidative stress markers (plasma F2-isoprostane) and alpha-tocopherol. The optimal time to significantly reduce plasma F2-isoprostane was determined from this study. In the dose response study, 60 patients randomly received placebo, placebo, 100, 200, 400, 800 or 1600 IU normal daily alpha-tocopherol (RRR) for the specified period of study. Blood was collected at baseline and every 2 weeks and analyzed for F2-isoprostane and alpha-tocopherol plasma. It is assumed that a dose of 800 IU of vitamin E is required to significantly reduce plasma F2 isoprostanes. This study determines the time and dose required for alpha-tocopherol to significantly reduce oxidative stress in hemodialysis patients. The data are used to evaluate the effect of alpha-tocopherol on cardiovascular outcomes in hemodialysis patients (Reed *et al.*, 2009).

Mode of Action

Failure of supplementary anti-oxidants to swap α -tocopherol in numerous cell-related responses, insufficient anti-oxidant rejoinder to expression of gene below deficit α -tocopherol, unrivaled acceptance of α -tocopherol and its leisureliel other catabolism reactions are actually prominent for the role of molecules of diverse tocopherols, that are pragmatic in-vitro. They are not disparate to their anti-oxidant possessions and all information actuality in sustenance of the non-antioxidant molecular purpose of α -tocopherol. Additionally, α -tocopherol is not enough at physiological

concentrations in order to defend contrary to oxidant-induced damage or avert illness purportedly produced by oxidative damage. α -tocopherol seems to work as a ligand of not yet recognized exact proteins (transcription factors and receptors) which are proficient to adapt gene expression and signal transduction (Azzi, 2007).

Reinvestigation of mechanism of action of α -tocopherol has been methodically accomplished in past few years. According to some new interventions, the chain-breaking antioxidant (α -tocopherol) is recognized as the main shield compared to lipid peroxidation (in vivo) (Ognjanovic *et al.*, 2003). α -tocopherol roles in active structures chiefly as a free radical scavenger and lipid anti-oxidant had to be reviewed (Ogru *et al.*, 2003). It has been verified that, between some significant detections which have carried to this novel standard is a verdict that of the 08 vitamin E (the homonymous tocotrienols and δ -, γ -, β -, α -tocopherol), solitary α -tocopherol seems to be engaged in momentous quantities by an organism (Wolf, 2006). This incident is actually a result of countenance of a protein (α -TTP) with augmented discrimination for α -tocopherol (Engin, 2009). The low or very low affinity for other tocopherols is not shared by the implicit message of specific evolutionary pressure by α -tocopherol, which is not equally powerful by other antioxidants. It is experimentally proved that α -tocopherol has a unique characteristic that it is capable to modify innumerable functions of cell in an exclusive mode that is not common in further anti-oxidants (Azzi *et al.*, 2000). Infinite observations specify that unlimited mechanisms of cell like cessation of platelet accumulation and union, cessation of cytokine proclamation and monocyte ROS, cessation of endothelial-monocyte union, endothelial veracity defense and cessation of smooth tissues propagation are all meticulous by the non-antioxidant possessions of α -tocopherol. (Ouazzani-Chahdi *et al.*, 2007). It is regarded as a much complicated but an acceptable directive of cellular roles that are arbitrated by non-governable permitted radical chain reactions (Janssens and Goris, 2001). Several further genes have been originated to be under the regulator of α -tocopherol after the innovative discovery that α -tocopherol is capable to control the expression of genes (Oommen *et al.*, 2007). Though, by an apparent compensatory mechanism, not a single gene expressed anti-oxidant enzymatic reaction in the absenteeism of α -tocopherol. Diverse tocotrienols and tocopherols have properties which are sovereign of their comparative anti-oxidant possessions at the level of cell. For example, diverse equivalents of carbonitrile byproducts and tocopherols constrain the propagation of smooth tissues by a systematic way which is distinct from anti-oxidant possessions of the particles (Gohil *et al.*, 2003). Furthermore, rivalry among β -tocopherol and α -tocopherol in constraining smooth muscle cell proliferation and PKC (Farbstein *et al.*, 2010) proposes the presence of necessary spot for the 02 particles and might not be clarified in relations of 02 anti-oxidants that are added together have fewer outcome as compared to α -tocopherol alone. α -Tocotrienol acts by an aid of parameter of gene expression in an anti-oxidant-free method (Rimbach and De Pascual-Teresa, 2005).

Adversative properties

As an alternative, the consequences of CVDs deterrence (HOPE) valuation appraised vitamin E and an angiotensin-converting enzyme inhibitor in people with high-risk cardiovascular events who had diabetes and at least one other cardiovascular disease. The results of vitamin E intervention had no significant benefits (Yusuf *et al.*, 2000). Negative results have also been reported in the cardio-protection Study, a placebo-controlled, double-blind, placebo-controlled, double-blind trial comparing antioxidant vitamin supplementation (600 mg vitamin E, 250 mg vitamin C, and 20 mg). The uses of Beta-carotene on daily basis with placebo cause approximately 20,000 high-risk cardiovascular patients. Although antioxidant treatment significantly increased blood vitamin D concentrations, it did not show beneficial effects in protecting cardiovascular disease. A meta-analysis involving 15 clinical trials on cardiovascular disease outcomes failed to show antioxidant benefits against

cardiovascular disease outcomes. This analysis included 8 studies of vitamin E (50 – 800 IU everyday) and most trials included > 1000 participants. Inappropriately, there have been some evidences of a lack of efficacy in different degrees and types of CVDs and in different populations of men and women (Vivekananthan *et al.*, 2003). According to the above data, primary and secondary prevention trials do not support the clinical effect of vitamin E as an antioxidant that reduces the burden of cardiovascular disease. Although vitamin E prevents lipid peroxidation, addition of vitamin E to lipid emulsions can accelerate lipid peroxidation in vitro (Becvarova *et al.*, 2005). In vivo studies have also shown that vitamin E administration increases plasma lipid peroxidation product levels in high-fat diet smokers and increased liver fat in mice fed a high-fat diet plus ethanol (Weinberg *et al.*, 2001). Adult vitamin E is generally thought to be of minimal toxicity, but blood coagulation is affected by vitamin K interference. In addition, a meta-analysis showed that high intake of vitamin E supplements may increase overall mortality in the general population. These results suggest that large amounts of vitamin E supplements can be harmful (Miller *et al.*, 2005).

Conclusion

Maximum of the risk factors of CVDs, counting diabetes, obesity, hypertension and hyperlipidemia are expressively pretentious with dietetic influences. Disproportionate eating of vegetables and fruits containing α -tocopherols has been linked with subordinate tolls of overweightness and thus a developed danger of 2 diabetes and hypertension. Considering the association among vegetable and fruit ingesting and cardiovascular disorders, it is mandatory to motivate cognizance of nutritional strategies and also customer tutoring in order to decrease the menace of CVDs. Solution is actually the intensification of entire ration of ≥ 05 regular mealtimes of vegetables and fruits. It must be eaten as per a basis of water, phytochemicals, fiber and vitamins as portion of a well-adjusted regime.

For optimum health and nutrition, people ought to attain bio-active nutrients from an extensive variety of entire diets or from those foods which are enrich in fiber or cereals derivative of vegetables but not from nutritional supplements. Carotene, vitamin C and vitamin E proved unsatisfactory in experimental prosecutions held earlier. This happened due to involvedness of available vegetables and fruits having a bulky quantity of biologically dynamic composites besides additional biologically dynamic foundations of biologically dynamic composites. As an alternative, indication recommends that a multifaceted customary of numerous nutritional composites interrelate to impact the menace of cardiovascular troubles. Consequently, consideration to entire nutritional outlines and nutriments is decisive to condense the menace of cardiovascular disease (CVD).

Evidence advocates that intake of vegetables and fruits have a defensive consequence contrary to cardiovascular disorders, principally from observational investigations. The true level of the association among vegetable and fruit ingesting and a prevalence of cardiovascular disorder are observed. It was not fully proved by the outcomes of controlled randomized prosecutions that consumption of vegetables and fruits preserve against cardiovascular disorders, partially as nutritional interferences have not been adequately concentrated to achieve an ideal investigation of their assumed belongings. Defensive mechanisms might comprise not only recognized organic nutrients like potassium and fiber but also practical characteristics of vegetable and fruit ingesting, such as energy density and low glycemic load. Collectively, it seemed that entire evidence congregated up to now assists an impression that augmented feasting of vegetables and fruits might decrease the menace of CVDs.

The consequence of quantity on the types of fruits and vegetables consumed the amount of processing in unpublished associations and an impact of certain classes of fruits and vegetables is

still undistinguishable. Few reservations might be addressed in imminent.

References

- Alissa, E. M., & Ferns, G. A. (2017). Dietary fruits and vegetables and cardiovascular diseases risk. *Critical reviews in food science and nutrition*, 57(9), 1950-1962.
- Azzi, A. (2007). Molecular mechanism of α -tocopherol action. *Free Radical Biology and Medicine*, 43(1), 16-21.
- Azzi, A., Breyer, I., Feher, M., Pastori, M., Ricciarelli, R., Spycher, S., ... & Zingg, J. M. (2000). Specific cellular responses to α -tocopherol. *The Journal of nutrition*, 130(7), 1649-1652.
- Babu, K. H. (2011). Reversal Of CCL4 Induced Toxicity In Rat Liver By Alfa Tocopherol.
- Becvarova, I., Saker, K. E., Swecker, W. S., & Troy, G. C. (2005). Peroxidative protection of parenteral admixture by D-alpha-tocopherol. *Veterinary Therapeutics*, 6(4), 280.
- Brandsma, E., Kloosterhuis, N. J., Koster, M., Dekker, D. C., Gijbels, M. J., van der Velden, S., ... & Fu, J. (2019). A proinflammatory gut microbiota increases systemic inflammation and accelerates atherosclerosis. *Circulation research*, 124(1), 94-100.
- Chen, C., Wang, Y., Cao, Y., Wang, Q., Anwaier, G., Zhang, Q., & Qi, R. (2019). Inhibitory effects and mechanism of probucol on elastase-induced abdominal aortic aneurysm in mice. *British Journal of Pharmacology*.
- Chepda, T., Cadau, M., Lassabliere, F., Reynaud, E., Perier, C., Frey, J., & Chamson, A. (2001). Synergy between ascorbate and α -tocopherol on fibroblasts in culture. *Life Sciences*, 69(14), 1587-1596.
- Dandapat, J., Chainy, G. B., & Rao, K. J. (2000). Dietary vitamin-E modulates antioxidant defence system in giant freshwater prawn, *Macrobrachium rosenbergii*. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 127(1), 101-115.
- Denova-Gutiérrez, E., Tucker, K. L., Salmerón, J., Flores, M., & Barquera, S. (2016). Relative validity of a food frequency questionnaire to identify dietary patterns in an adult Mexican population. *salud pública de méxico*, 58, 608-616.
- Dietrich, M., Traber, M. G., Jacques, P. F., Cross, C. E., Hu, Y., & Block, G. (2006). Does γ -tocopherol play a role in the primary prevention of heart disease and cancer? A review. *Journal of the American College of Nutrition*, 25(4), 292-299.
- Doria-Lamba, L., De Grandis, E., Cristiani, E., Fiocchi, I., Montaldi, L., Grosso, P., & Gellera, C. (2006). Efficacious vitamin E treatment in a child with ataxia with isolated vitamin E deficiency. *European journal of pediatrics*, 165(7), 494-495
- DRI. (2005). Institute of Medicine, Food and Nutrition Board, Dietary Reference Intakes: energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids.

- Engin, K. N. (2009). Alpha-tocopherol: looking beyond an antioxidant. *Molecular vision*, 15, 855.
- Falk, J., Krauß, N., Dähnhardt, D., & Krupinska, K. (2002). The senescence associated gene of barley encoding 4-hydroxyphenylpyruvate dioxygenase is expressed during oxidative stress. *Journal of Plant Physiology*, 159(11), 1245-1253.
- Farbstein, D., Kozak-Blickstein, A., & Levy, A. P. (2010). Antioxidant vitamins and their use in preventing cardiovascular disease. *Molecules*, 15(11), 8098-8110.
- Fujisawa, S., Atsumi, T., Ishihara, M., & Kadoma, Y. (2004). Cytotoxicity, ROS-generation activity and radical-scavenging activity of curcumin and related compounds. *Anticancer research*, 24(2B), 563-570.
- Gale, C. R., Ashurst, H. E., Powers, H. J., & Martyn, C. N. (2001). Antioxidant vitamin status and carotid atherosclerosis in the elderly. *The American journal of clinical nutrition*, 74(3), 402-408.
- Ganapathi, A., Roy, A., & Savitha, G. (2018). Estimation of C-Reactive Protein in Cardiovascular Disease Patients. *Research Journal of Pharmacy and Technology*, 11(8), 3303-3307.
- Gohil, K., Schock, B. C., Chakraborty, A. A., Terasawa, Y., Raber, J., Farese Jr, R. V., ... & Traber, M. G. (2003). Gene expression profile of oxidant stress and neurodegeneration in transgenic mice deficient in α -tocopherol transfer protein. *Free Radical Biology and Medicine*, 35(11), 1343-1354.
- Gong, C., Ning, P., & Bai, J. (2012). Responses of antioxidative protection to varying drought stresses induced by micro-ecological fields on desert C4 and C3 plants in Northwest China. *Life Sci. J*, 9(4), 2006-2016.
- Grammas, P., Hamdheydari, L., Benaksas, E. J., Mou, S., Pye, Q. N., Wechter, W. J., ... & Hensley, K. (2004). Anti-inflammatory effects of tocopherol metabolites. *Biochemical and biophysical research communications*, 319(3), 1047-1052.
- Graves, D. B., & Bauer, G. (2018). Key roles of reactive oxygen and nitrogen species. In *Comprehensive clinical plasma medicine* (pp. 71-82). Springer, Cham.
- Hijmans, J. G., Bammert, T. D., Stockelman, K. A., Reiakvam, W. R., Greiner, J. J., & DeSouza, C. A. (2019). High glucose-induced endothelial microparticles increase adhesion molecule expression on endothelial cells. *Diabetology international*, 10(2), 143-147.
- Jain, P. K., Sharma, P., & Joshi, S. C. (2019). Antioxidant And Lipid Lowering Effects Of *Elaeocarpus Ganitrus* In Cholesterol Fed Rabbits. *International Journal Of Pharmaceutical Sciences And Research*, 9(2), 526-534.
- Janssens, V., & Goris, J. (2001). Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. *Biochemical Journal*, 353(Pt 3), 417.
- Jargin, S. V. (2018). The potential use of flavonoids as venoactive drugs and the role of citrus

fruits. *Journal Of Intercultural Ethnopharmacology*, 7(1), 97-100.

Kang, J. H., Cook, N., Manson, J., Buring, J. E., & Grodstein, F. (2006). A randomized trial of vitamin E supplementation and cognitive function in women. *Archives of internal medicine*, 166(22), 2462-2468.

Kim, J. Y., Kim, M. S., Jeong, S., & Kwon, O. (2018). New paradigm for human intervention study in functional food development. *Food Science and Industry*, 51(1), 8-15.

Lim, Y., & Traber, M. G. (2007). Alpha-tocopherol transfer protein (α -TTP): Insights from alpha-tocopherol transfer protein knockout mice. *Nutrition research and practice*, 1(4), 247-253.

Liu, H., Peng, H., Xiang, H., Guo, L., Chen, R., Zhao, S., ... & Chen, S. (2018). TWEAK/Fn14 promotes oxidative stress through AMPK/PGC-1 α /MnSOD signaling pathway in endothelial cells. *Molecular medicine reports*, 17(1), 1998-2004.

Malekmohammad, K., Sewell, R. D., & Rafeian-Kopaei, M. (2019). Antioxidants and Atherosclerosis: Mechanistic Aspects. *Biomolecules*, 9(8), 301.

Manivannan, P., Jaleel, C. A., Kishorekumar, A., Sankar, B., Somasundaram, R., Sridharan, R., & Panneerselvam, R. (2007). Changes in antioxidant metabolism of *Vigna unguiculata* (L.) Walp. by propiconazole under water deficit stress. *Colloids and Surfaces B: Biointerfaces*, 57(1), 69-74.

Martínez, J. P., Silva, H. F. L. J., Ledent, J. F., & Pinto, M. (2007). Effect of drought stress on the osmotic adjustment, cell wall elasticity and cell volume of six cultivars of common beans (*Phaseolus vulgaris* L.). *European Journal of Agronomy*, 26(1), 30-38.

Massacci, A., Nabiev, S. M., Pietrosanti, L., Nematov, S. K., Chernikova, T. N., Thor, K., & Leipner, J. (2008). Response of the photosynthetic apparatus of cotton (*Gossypium hirsutum*) to the onset of drought stress under field conditions studied by gas-exchange analysis and chlorophyll fluorescence imaging. *Plant Physiology and Biochemistry*, 46(2), 189-195.

Mead, A., Atkinson, G., Albin, D., Alphey, D., Baic, S., Boyd, O., ... & Greene, P. (2006). Dietetic guidelines on food and nutrition in the secondary prevention of cardiovascular disease—evidence from systematic reviews of randomized controlled trials (second update, January 2006). *Journal of human nutrition and dietetics*, 19(6), 401-419.

Miller, E. R., Pastor-Barriuso, R., Dalal, D., Riemersma, R. A., Appel, L. J., & Guallar, E. (2005). Meta-analysis: high-dosage vitamin E supplementation may increase all-cause mortality. *Annals of internal medicine*, 142(1), 37-46.

Miyoshi, N., Naniwa, K., Kumagai, T., Uchida, K., Osawa, T., & Nakamura, Y. (2005). α -Tocopherol-mediated caspase-3 up-regulation enhances susceptibility to apoptotic stimuli. *Biochemical and biophysical research communications*, 334(2), 466-473.

Munné-Bosch, S. (2005). The role of α -tocopherol in plant stress tolerance. *Journal of plant physiology*, 162(7), 743-748.

Nadeem, N., Woodside, J. V., Kelly, S., Allister, R., Young, I. S., & McEneny, J. (2012). The two faces of α - and γ -tocopherols: an in vitro and ex vivo investigation into VLDL, LDL and HDL oxidation. *The Journal of nutritional biochemistry*, 23(7), 845-851.

Nordberg, J., & Arnér, E. S. (2001). Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free radical biology and medicine*, 31(11), 1287-1312.

Ognjanovic, B. I., Pavlovic, S. Z., Maletic, S. D., Zikic, R. V., Stajn, A. S., Radojicic, R. M., ... & Petrovic, V. M. (2003). Protective influence of vitamin E on antioxidant defense system in the blood of rats treated with cadmium. *Physiological Research*, 52(5), 563-570.

Ogru, E., Gianello, R., Libinaki, R., Smallridge, A., Bak, R., Geytenbeek, S., ... & West, S. (2003). Vitamin E phosphate: an endogenous form of vitamin E. *Medimond Srl*, 127-132.

Ohm, V. A., Stöckmann, H., & Schwarz, K. (2005). The more—The better? Estimating the inhibitory activity of α -tocopherol towards lipid oxidation. *Journal of plant physiology*, 162(7), 785-789.

Oommen, S., Vasu, V. T., Leonard, S. W., Traber, M. G., Cross, C. E., & Gohil, K. (2007). Genome wide responses of murine lungs to dietary α -tocopherol. *Free radical research*, 41(1), 98-133.
Ouazzani-Chahdi, A., Elimadi, A., Chabli, A., Spenard, J., Colin, P., & Haddad, P. S. (2007). Combining ursodeoxycholic acid or its NO-releasing derivative NCX-1000 with lipophilic antioxidants better protects mouse hepatocytes against amiodarone toxicity. *Canadian journal of physiology and pharmacology*, 85(2), 233-242.

Parthasarathy, S., Raghavamenon, A., Garelnabi, M. O., & Santanam, N. (2010). Oxidized low-density lipoprotein. In *Free Radicals and Antioxidant Protocols* (pp. 403-417). Humana Press.

Ranjha, M. M. A. N., Amjad, S., Ashraf, S., Khawar, L., Safdar, M. N., Jabbar, S., ... & Murtaza, M. A. (2020a). Extraction of Polyphenols from Apple and Pomegranate Peels Employing Different Extraction Techniques for the Development of Functional Date Bars. *International Journal of Fruit Science*, DOI: 10.1080/15538362.2020.1782804

Ranjha, M. M. A. N., Irfan, S., Nadeem, M., & Mahmood, S. (2020b). A Comprehensive Review on Nutritional Value, Medicinal Uses, and Processing of Banana. *Food Reviews International*, DOI: 10.1080/87559129.2020.1725890

Raitakari, O. T., Juonala, M., Kähönen, M., Taittonen, L., Laitinen, T., Mäki-Torkko, N., ... & Åkerblom, H. K. (2003). Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *Jama*, 290(17), 2277-2283.

Reed, A., Cho, Y. J., Coombes, J. S., & Fassett, R. G. (2009). Time course and dose response of α -tocopherol on oxidative stress in haemodialysis patients. *BMC nephrology*, 10(1), 32.

Refa, S., Wardhani, C. K., & Dewi, N. A. (2019). The effect of lutein and zeaxanthin administration on reactive oxygen species levels in blood serum and contrast sensitivity changes in patients with non-proliferative diabetic retinopathy. *Drug Invention Today*, 11(10).

Rimbach, G., & De Pascual-Teresa, S. (2005). Application of nutrigenomics tools to analyze the role of oxidants and antioxidants in gene expression. *OXIDATIVE STRESS AND DISEASE*, 17, 1.

Rizvi, S., Raza, S. T., Ahmed, F., Ahmad, A., Abbas, S., & Mahdi, F. (2014). The role of vitamin E in human health and some diseases. *Sultan Qaboos University Medical Journal*, 14(2), e157.

Ryan, E., Galvin, K., O'Connor, T. P., Maguire, A. R., & O'Brien, N. M. (2007). Phytosterol, squalene, tocopherol content and fatty acid profile of selected seeds, grains, and legumes. *Plant Foods for Human Nutrition*, 62(3), 85-91.

Satija, A., Yu, E., Willett, W. C., & Hu, F. B. (2015). Understanding nutritional epidemiology and its role in policy. *Advances in nutrition*, 6(1), 5-18.

Schaich, K. (2006). 24 Lipid Oxidation in Specialty Oils. *Nutraceutical and specialty lipids and their co-products*, 401.

Sen, C. K., Khanna, S., & Roy, S. (2006). Tocotrienols: Vitamin E beyond tocopherols. *Life sciences*, 78(18), 2088-2098.

Serin, O., Konukoglu, D., Firtina, S., & Mavis, O. (2007). Serum oxidized low density lipoprotein, paraoxonase 1 and lipid peroxidation levels during oral glucose tolerance test. *Hormone and metabolic research*, 39(03), 207-211.

Shils, M. E., & Shike, M. (Eds.). (2006). *Modern nutrition in health and disease*. Lippincott Williams & Wilkins.

Silva, E. H., Wickramatilake, C. M., Lekamwasam, S., Mudduwa, B., Kumari, L., & Ubayasiri, R. A. (2019). Vascular dysfunction and atherosclerosis in chronic kidney disease; A distinct entity. *Journal of Nephropathology*, 8(2).

Singh, A., Gowtham, S., Chakrapani, L. N., Ashokkumar, S., Kumar, S. K., Prema, V., ... & Sathyamoorthy, Y. K. (2018). Aegeline vs Statin in the treatment of Hypercholesterolemia: A comprehensive study in rat model of liver steatosis. *Functional Foods in Health and Disease*, 8(1), 1-16.

Sivaraman, D., Anbu, N., Kabilan, N., Kumar, M. P., Shanmugapriya, P., & Christian, G. J. (2019). REVIEW ON CURRENT TREATMENT STRATEGY IN ALZHEIMER'S DISEASE AND ROLE OF HERBS IN TREATING NEUROLOGICAL DISORDERS. *Int J Trans Res Ind Med*, 1(1), 33-43.

Sun, Y., Nemec-Bakk, A. S., Mallik, A. U., Bagchi, A. K., Singal, P. K., & Khaper, N. (2019). Blueberry extract attenuates doxorubicin-induced damage in H9c2 cardiac cells. *Canadian journal of physiology and pharmacology*, 97(9), 880-884.

Tegischer, K., Tausz, M., Wieser, G., & Grill, D. (2002). Tree-and needle-age-dependent variations in antioxidants and photoprotective pigments in Norway spruce needles at the alpine timberline. *Tree physiology*, 22(8), 591-596.

Traber, M. G. (2007). Vitamin E regulatory mechanisms. *Annu. Rev. Nutr.*, 27, 347-362.

- Traber, M. G., & Atkinson, J. (2007). Vitamin E, antioxidant and nothing more. *Free radical biology and medicine*, 43(1), 4-15.
- Tucker, J. M., & Townsend, D. M. (2005). Alpha-tocopherol: roles in prevention and therapy of human disease. *Biomedicine & pharmacotherapy*, 59(7), 380-387.
- Ulatowski, L. M., & Manor, D. (2015). Vitamin E and neurodegeneration. *Neurobiology of disease*, 84, 78-83.
- Ulatowski, L., Dreussi, C., Noy, N., Barnholtz-Sloan, J., Klein, E., & Manor, D. (2012). Expression of the α -tocopherol transfer protein gene is regulated by oxidative stress and common single-nucleotide polymorphisms. *Free Radical Biology and Medicine*, 53(12), 2318-2326.
- Vasiliki, G., Charalampia, D., & Haralabos, K. C. (2019). In vitro antioxidant, antithrombotic, antiatherogenic and antidiabetic activities of *Urtica dioica*, *Sideritis euboica* and *Cistus creticus* water extracts and investigation of pasta fortification with the most bioactive one. *Current pharmaceutical biotechnology*.
- Vivekananthan, D. P., Penn, M. S., Sapp, S. K., Hsu, A., & Topol, E. J. (2003). Use of antioxidant vitamins for the prevention of cardiovascular disease: meta-analysis of randomised trials. *The Lancet*, 361(9374), 2017-2023.
- Weinberg, R. B., VanderWerken, B. S., Anderson, R. A., Stegner, J. E., & Thomas, M. J. (2001). Pro-oxidant effect of vitamin E in cigarette smokers consuming a high polyunsaturated fat diet. *Arteriosclerosis, thrombosis, and vascular biology*, 21(6), 1029-1033.
- Windisch, M. (2000). Approach towards an integrative drug treatment of Alzheimer's disease. In *Advances in Dementia Research* (pp. 301-313). Springer, Vienna.
- Wolf, G. (2006). How an increased intake of alpha-tocopherol can suppress the bioavailability of gamma-tocopherol. *Nutrition reviews*, 64(6), 295-299.
- Yoshida, H. (2018). An Intriguing and Important Concept Relevant to Oxidized Low-Density Lipoprotein and Atherogenesis is Still Problematic for its Contribution to the Better Understanding of Clinical Atherosclerosis. *Journal of atherosclerosis and thrombosis*, 25(10), 1007-1008.
- Yusuf, S., Dagenais, G., Pogue, J., Bosch, J., & Sleight, P. (2000). Vitamin E supplementation and cardiovascular events in high-risk patients. *The New England journal of medicine*, 342(3), 154-160.
- Zhang, X., & Schmidt, R. E. (2000). Hormone-containing products' impact on antioxidant status of tall fescue and creeping bentgrass subjected to drought. *Crop Science*, 40(5), 1344-1349.
- Zhu, Y., Anand, R., Geng, X., & Ding, Y. (2018). A mini review: garlic extract and vascular diseases. *Neurological research*, 40(6), 421-425.
- Zhu, Y., Xian, X., Wang, Z., Bi, Y., Chen, Q., Han, X., & Chen, R. (2018). Research progress on the relationship between atherosclerosis and inflammation. *Biomolecules*, 8(3), 80.

Zou, C. G., Agar, N. S., & Jones, G. L. (2001). Oxidative insult to human red blood cells induced by free radical initiator AAPH and its inhibition by a commercial antioxidant mixture. *Life Sciences*, 69(1), 75-86.



© 2019 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format. Adapt — remix, transform, and build upon the material for any purpose, even commercially. The licensor cannot revoke these freedoms as long as you follow the license terms. Under the following terms: Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use. No additional restrictions. You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits