



History:

Received: February 4, 2019
Reviewed: May 31, 2019
Accepted: June 3, 2019
First Published: June 17, 2019
Collection year: 2019
Status: Published

Identifiers and Pagination:

Year: 2019
Volume: 4
First Page: 1
Last Page: 7
Publisher Id: AdvFood&NutrSci.4.1
DOI:<http://dx.doi.org/10.21065/AdvFoodNutriSci.4.1>

Corresponding author:

Dr. Nelson A Cox, 950 College Station
Road Athens, GA 30605, T.: (706)546-
3484 (phone), F.: (706)546-3633, E.:
nelson.cox@ars.usda.gov

Citation:

Landrum MA, NA Cox, JL Wilson, ME
Berrang, GR Gamble, MA Harrison, BD
Fairchild, WK Kim, and A Hinton, Jr.
Reduction of campylobacter on poultry
thighs using sequential treatments of
antimicrobials. Adv Food & Nutr Sci.
Vol. 4. 2019. p 1-7

Original Research Article

REDUCTION OF *CAMPYLOBACTER* ON POULTRY THIGHS USING SEQUENTIAL TREATMENTS OF ANTIMICROBIALS

Landrum MA¹, NA Cox², JL Wilson¹, ME Berrang², GR Gamble², MA Harrison¹, BD Fairchild¹, WK Kim¹, and A Hinton, Jr²

1. University of Georgia, Athens, GA;
2. The U.S. National Poultry Research Center, Athens, GA

Running Title: Sequential Dips to Reduce *Campylobacter*

Keywords: *Campylobacter*, CMS PoultryHresh™, broiler, carcass rinse, immersion

Abstract

Campylobacter is a major concern for poultry processors, as USDA performance standards have become stricter. This study evaluated CMS PoultryHresh™, a low pH processing aid, and peracetic acid using consecutive and sequential dip treatments to reduce *Campylobacter* in thighs. Thighs (n=3/treatment group) were inoculated with a *C. coli* marker strain (10⁸) and each dipped into bags containing 1 L of treatment 1 for 6 s. Thighs were allowed 5 s to drip, placed onto foil for 60 s, and dipped into treatment 2 for 6 s. After 5 s drip time, each was placed in a bag with 150 mL buffered peptone water and hand shaken for 60 s; controls same procedure, no treatment. Rinsates were serially diluted, plated onto Campy Cefex agar with 200 ppm gentamicin and incubated microaerobically for 48 h at 42°C. Procedures were replicated 5 times. Significant reductions compared to untreated using consecutive dips of PoultryHresh™ and PAA were 98.2% and 99.3%, respectively. Treatments of PoultryHresh™ then peracetic acid reduced *Campylobacter* 99.2% from untreated thighs. Peracetic acid then PoultryHresh™ showed significant reductions compared to all other treatments (99.9% from untreated). Treating with this sequence may allow processors to meet the strict performance standards on *Campylobacter* in broiler parts.

Introduction

Campylobacter is the main cause of bacterial gastroenteritis in the world and researchers have shown it to be present in high levels in retail poultry. [1],[2] Of *Campylobacter* illnesses, 50 - 70% were caused by consumption of poultry or poultry products. [3],[4] Symptoms of *Campylobacter* include fever, abdominal pain and diarrhea within 2 - 5 days of ingesting contaminated product. [5],[6] Stricter performance standards have been implemented by the USDA, aimed at reducing incidence of *Campylobacter* in processing facilities. [7] FSIS estimated 46% of poultry facilities would not be able to meet new requirements; therefore, processors must give serious attention to finding efficient antimicrobial treatments to be used throughout the plant. It is mandatory *Campylobacter* prevalence remain below 7.7% on poultry parts (4 of 52 samples), allowing only a small margin of error. [7]

A variety of antimicrobials have been evaluated for use on poultry at various points

Funding:

The authors received no direct funding for this research.

Competing Interests:

The authors declare no competing interests

Additional information is available at the end of the article.

throughout processing procedures. It is essential chemicals be cost effective, reduce pathogen prevalence and not cause organoleptic damage to poultry carcasses or parts. [8] Antimicrobials vary in levels of efficacy, treatment concentration, contact time and application method. Chemical efficacy is affected by microbial load, composition of flora, organic material or possibly changed from residual effects of prior treatments. [9] Stopforth et al. [10] determined a single intervention could not significantly reduce pathogen presence on finished carcasses. Instead, a multi-hurdle approach is required at various intervention points throughout processing for increased pathogen reduction on poultry products. [10],[11] There was no previous research found evaluating consecutive, sequential treatments and how they affect pathogens. The purpose of this study is to evaluate the effectiveness of a low acid antimicrobial, CMS PoultrypHresh™, and whether dipping carcasses in consecutive or sequential treatments along with peracetic acid (PAA) leads to further reductions in the prevalence of *Campylobacter* on poultry thighs.

Materials and Methods

Bacterial Marker Strain

Cox et al. [12] developed a marker strain of *Campylobacter* resistant to the antibiotic gentamicin, *Campylobacter coli* (CC^{GR}), which was used in this study. Campy Cefex agar (Sigma, St. Louis, MO) plates were prepared using 200 ppm gentamicin (CCGen) to eliminate the growth of wild type *Campylobacter* that may be naturally present on poultry thighs. Initial cultures were streaked from storage in an -80°C freezer, maintained in Bolton's broth with 15% glycerol and no supplements. Cultures were streaked onto CCGen agar and incubated in sealed bags under microaerobic (5% O₂, 10% CO₂, 85% N₂) conditions at 42°C for 48 h. A sterile swab was used to remove colonies and mix them into a 9 mL tube of Bolton's broth, placed in sealed bags and incubated microaerobically for 24 h at 42°C. Tubes were evaluated for colorimeter analysis by a Spectronic 200E (Thermo Fisher Scientific, Madison, WI) and approximately a 10⁸ cfu/mL CC^{GR} inoculum was prepared. Inocula were confirmed using serial dilutions, plated on CCGen and incubated for 48 h at 42°C.

Treatment Procedures

Skin-on poultry thighs were purchased from a local grocery (n=18) and divided into the six treatment groups. A 6.5 cm² section of skin was inoculated on each thigh by spreading 0.1 mL of 10⁸ CC^{GR} marker strain. Each thigh was inoculated individually and given a 5 min attachment period before treatment. Thighs were individually dipped into Ziploc bags containing either 1 liter of water, PAA (600 ppm), or PoultrypHresh (pH 1.5) for 6 s. Thighs were removed and placed on tin foil squares for 60 s. Each thigh was then dipped into the second dip treatment for 6 s, drained 5 s, placed into individual sealable bags with 150 mL buffered peptone water (BPW), and hand shaken and massaged for 60 s. Thighs were removed, and rinses placed on ice for approximately 15 min before diluting. Three thighs served as controls in each replication and were inoculated as described, remained untreated and placed directly into rinse bags with BPW after the attachment period. The entire experiment was replicated 5 times.

Plating and Incubation

Rinsates were serially diluted and a plate spreader used to disperse onto CCGen agar plates in duplicate. Plates were incubated at 42°C for 48 h. Characteristic colonies were counted, and cfu/mL log transformed.

Statistical Analysis

The study was constructed of five replicates of skin-on thighs using 18 individual thighs (N=18; n=3). Each replicate consisted of 3 of each untreated, consecutive water dips, consecutive PoultrypHresh™ (pH 1.5) dips, consecutive PAA (600 ppm) dips, PoultrypHresh™ followed by PAA dip, and PAA followed by PoultrypHresh™ dip. Duplicate counts were averaged, and numbers were transformed by log₁₀. Data was analyzed using Statistica software (Statistica, 2013). A General Linear Model was conducted to determine whether sequential dips were statistically different ($P < 0.05$). Means were separated with a Tukey Multiple Comparison test; statistical significance was assigned at $P \leq 0.05$.

RESULTS

Consecutive water dips were used to determine rinsing effect of water alone. A 0.8 log₁₀ cfu/mL *Campylobacter* reduction was observed, although contamination remained high at 4.9 log₁₀ cfu/mL (Table 1). Treating with consecutive PoultrypHresh™ dips significantly ($P \leq 0.05$) reduced *Campylobacter* from 5.64 log₁₀ cfu/mL untreated and 4.87 log₁₀ cfu/mL water dipped to 3.90 log₁₀ cfu/mL (Table 1). This equates to a reduction of 98.2% from untreated and 89.3% from water dipped thighs (Table 1). Dipping with consecutive PAA dips showed slightly higher reductions; reducing *Campylobacter* 99.3% (2.1 log₁₀ cfu/mL) from untreated and 95.7% (1.4 log₁₀ cfu/mL) from water dipped samples (Table 1).

Dipping in PoultrypHresh™ followed by PAA demonstrated findings similar to consecutive PAA dips. Interestingly, thighs dipped in PAA followed by PoultrypHresh™, showed reductions significantly ($P \leq 0.05$) greater than any other dipping combination. *Campylobacter* was reduced 99.9% (> 3 log₁₀ cfu/mL reduction) from untreated thighs and 99.6% (2.4 log₁₀ cfu/mL reduction) from water treated (Table 1). This pattern was observed in all 5 replications of the study; therefore, the order of chemicals used in dipping sequences was significantly ($P \leq 0.05$) correlated to *Campylobacter* reductions.

DISCUSSION

Results can be compared to a study by Landrum et al. (2018) evaluating PoultrypHresh™ on broiler thighs, where *Campylobacter* reductions were higher compared to untreated from this study, 2.2 log₁₀ cfu/mL and 1.7 log₁₀ cfu/mL, respectively. Comparing the water dipped thighs to untreated, the research by Landrum et al. (2018) demonstrated a 1.4 log₁₀ cfu/mL reduction, whereas this study a 1.0 log₁₀ cfu/mL. Differences between the two studies were the dip time, (25 s compared to two 6 s dips) and air agitation, not used in this study. Shorter dip times of 6 s were chosen for this research as a more practical time of exposure in a modern processing facility. The higher dip times used by Landrum et al. (2018) exhibited better results, seemingly due to the longer exposure time, although the effects of air agitation could have also assisted in higher reductions. The improved microbial

reductions using air agitation have been shown in previous research. [14] Therefore, air agitation is a concept that may require more research to heighten reductions of microbial contamination.

PAA is a low pH organic peroxide mixture of acetic acid and hydrogen peroxide currently being used throughout the United States in multiple processing facilities. [8],[15],[16] The mode of action for PAA is a disruption to the cell membrane permeability, altering protein synthesis, leading to bacterial death. [17] PAA reduced the presence of *Campylobacter* slightly more than consecutive PoultrypHresh™ dips, although differences were not significant. Findings by Bauermeister et al. [8] showed a 1.5 log₁₀ cfu/mL *Campylobacter* reduction using an extremely low concentration of only 200 ppm PAA. As PAA is approved for use throughout processing at concentrations up to 2000 ppm, reductions could be even greater with increased acid levels. [18] Bauermeister [19] used 200 ppm PAA, only demonstrating a 1.5 log₁₀ cfu/mL reduction; therefore, other factors could be associated to its efficacy. King et al. [20] determined that the effects of PAA may vary and greatly depend on bacteria level and how they are attached to the surface. Nagel et al. [21] evaluated higher levels of PAA at 400 and 1000 ppm and reduced *Campylobacter* levels by more than 2.0 log₁₀ cfu/mL.

Scientists have determined a multi-hurdle approach is necessary for adequate pathogen reduction, which uses multiple intervention points throughout processing procedures such as the inside-outside bird washer, brush washer, cabinet washer, or dip tank before and/or after chilling. [22] Using this approach, the facility does not rely only on a single step intervention, but instead incorporates many applications for reducing foodborne pathogen prevalence prior to entering secondary processing [10],[23],[24] No research was found, however, demonstrating the usefulness of consecutive, sequential chemical hurdles for pathogen reduction. This study demonstrates the possibility that consecutive application of chemicals in a specific sequential order may reduce *Campylobacter* prevalence.

Table 1. Average Log₁₀ cfu/mL of *Campylobacter coli* recovered by replicate from sequentially dip treated thighs with no treatment, water-water, PpH-PpH, PAA-PAA, PpH-PAA, or PAA-PpH (mean ± standard error).

Treatment		Average Log ₁₀ cfu/mL	Reduction compared to Untreated (%)	Reduction compared to Water (%)
1 st Dip	2 nd Dip			
Untreated		5.64 ^a ± 0.04	-	-
Water	Water	4.87 ^b ± 0.05	83.0	-
PpH ¹	PpH	3.90 ^c ± 0.06	98.2	89.3
PAA ²	PAA	3.50 ^d ± 0.08	99.3	95.7
PpH	PAA	3.53 ^d ± 0.08	99.2	95.4
PAA	PpH	2.52 ^e ± 0.09	99.9	99.6

¹ PpH = CMS PoultrypHresh™

² PAA = peracetic acid

CONCLUSIONS

Findings from this study demonstrated PoultryHresh™ and PAA could reduce the prevalence of *Campylobacter* on poultry parts when used separately. However, when evaluated sequentially, PAA followed by PoultryHresh™ can significantly reduce pathogen presence. Such findings may be extremely beneficial and lead to better intervention strategies in future antimicrobial applications.

REFERENCES

1. Harrison WA, Griffith CJ, Tennant D, Peters AC. Incidence of *Campylobacter* and *Salmonella* isolated from retail chicken and associated packaging in South Wales. *Lett Appl Microbiol* 2001; 33:450-454.
2. Moore JE, Wilson TS, Wareing DR, Humphrey TJ, Murphy PG. Prevalence of thermophilic *Campylobacter* spp. in ready-to-eat foods and raw poultry in Northern Ireland. *J Food Prot* 2002; 65:1326-1328.
3. Allos B. *Campylobacter jejuni* infections: update on emerging issues and trends. *Clin Infectious Dis* 2001; 32(8):1201-1206.
4. Humphrey T, O'Brien S, Madsen M. *Campylobacters* as zoonotic pathogens: a food production perspective. *Int J Food Microbiol* 2007; 117:237-257.
5. Robinson DA. Infective dose of *Campylobacter jejuni* in milk. *Br. Med. J.* 1981; 282:1584.
6. Black RE, Levine MM, Clements ML, Hughs TP, Blaser MJ. Experimental *Campylobacter jejuni* infection in humans. *J Infect Dis* 1988; 157:472-479.
7. Food Safety and Inspection Service (FSIS). Pathogen Reduction - *Salmonella* and *Campylobacter* Performance Standards Verification Testing. United States Department of Agriculture. [Online] 2015. [cited 2018]. Available at: https://www.fsis.usda.gov/wps/wcm/connect/b0790997-2e74-48bf-9799-85814bac9ceb/28_IM_PR_Sal_Campy.pdf?MOD=AJPEREShttps://www.fsis.usda.gov/wps/wcm/connect/b0790997-2e74-48bf-9799-85814bac9ceb/28_IM_PR_Sal_Campy.pdf?MOD=AJPERES
8. Bauermeister LJ., Bowers JW, Townsend JC, McKee SR. Validating the efficacy of peracetic acid mixture as an antimicrobial in poultry chillers. *J Food Prot* 2008; 71:1119-1122.
9. Gill CO, Badoni M. Effects of peroxyacetic acid, acidified sodium chlorite or lactic acid solutions on the microflora of chilled beef carcasses. *Int J Food Microbiol* 2004; 91:43-50.
10. Stopforth JD, O'Connor R, Lopes M, Kottapalli B, Hill WE, Samadpour M. Validation of individual and multiple-sequential interventions for reduction of microbial populations during processing of poultry carcasses and parts. *J Food Prot* 2007; 70:1393-1401.
11. Bacon RT, Belk KE, Sofos JN, Clayton RP, Reagan JO, Smith GC. Microbial populations on animal hides and beef carcasses at different stages of slaughter in plants employing multiple-sequential interventions for decontamination. *J Food Prot* 2000; 63:1080-1086.
12. Cox NA, Richardson LJ, Berrang ME, Fedorka-Cray PJ, Buhr RJ. *Campylobacter coli* naturally resistant to elevated levels of gentamicin as a marker strain in poultry

- research. J Food Prot 2009; 72:1288-1292.
13. Landrum MA, Cox NA, Cosby DE, Berrang ME, Hinton A Jr., Mize SC, Jackson JS. Reduction of *Campylobacter* on chicken livers using a low acid processing aid. Adv Food Nutr Sci 2018; 3:1-6.
14. Bennett P. 2008. Compliance guidelines for controlling *Salmonella* and *Campylobacter* in poultry. 2nd ed. [Online] 2008 [cited 2018]. Available at: http://www.fsis.usda.gov/wps/wcm/connect/1d562776-ebfb4ea5-a19b-e994776a02c6/Compliance_Guideline_Controlling_Salmonella_Poultry.pdf?MOD=AJPERES.
15. USDA-FSIS. Sanitation performance standards compliance guide. Title 9. Code of Federal Regulations 416. [Online] 2016 [cited 2018]. Available at: <https://www.fsis.usda.gov/wps/portal/fsis/topics/regulatory-compliance/compliance-guides-index/sanitation-performance-standards/sanitation-compliance-guide>
16. [16] Sukted N, Tuitemwong P, Tuitemwong K, Poonlapdech W, Erickson LE. Inactivation of *Campylobacter* during immersion chilling of chicken carcasses. J Food Eng 2017; 202:25-33.
17. [17] Oyarzabal OA. Reduction of *Campylobacter* spp. by commercial antimicrobials applied during the processing of broiler chickens: a review from the United States perspective. J Food Prot 2005; 68(8):1752-1760.
18. [18] FSIS Directive. Safe and suitable ingredients used in the production of meat, poultry, and egg products. FSIS Directive 7120.1 Rev. 46. [Online] 2018 [cited 2018]. Available at: <https://www.fsis.usda.gov/wps/wcm/connect/bab10e09-ae4a-483b-8be8-809a1f051d4c/7120.1.pdf?MOD=AJPERES>
19. Bauermeister LJ. Evaluation of poultry meat safety and quality using peracetic acid in poultry chillers. Dissertation Auburn University. [Online] 2015 [cited 2018]. Available at: https://etd.auburn.edu/bitstream/handle/10415/4959/Laura_J_Bauermeister_Dissertation_12.12.2015.pdf;sequence=2
20. King DA, Lucia LM, Castillo A, Acuff GR, Harris KB, Savell JW. Evaluation of peracetic acid as a post-chilling intervention for control of *Escherichia coli* 0157:H7 and *Salmonella* Typhimurium on beef carcass surfaces. Meat Sci 2005; 69:401-407.
21. Nagel GM, Bauermeister LJ, Bratcher CL, Singh M, McKee SR. *Salmonella* and *Campylobacter* reduction and quality characteristics of poultry carcasses treated with various antimicrobials in a post-chill immersion tank. Int J Food Prot 2013; 165:281-286.
22. Smith J, Corkran S, McKee SR, Bilgili SF, Singh M. Evaluation of post-chill applications of antimicrobials against *Campylobacter jejuni* on poultry carcasses. J. Appl. Poult. Res. 2015; 24:451-456.
23. Zweifel C, Stephan R. Microbial Decontamination in the Food Industry: Microbial Decontamination of Poultry Carcasses. Woodhead Publishing. 2012; 60-95.
24. Sofos JN, Flick G, Nychas G, O'Bryan CA, Ricke SC, Crandall PG. Food Microbiology: Fundamentals and Frontiers. ASM Press. 2013; 111-167.



© 2017 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits